


CERTICAN.®

Wirkt und schützt – von Anfang an¹⁻⁸



1. Tedesco-Silva H et al. Everolimus Plus Reduced-Exposure CsA versus Mycophenolic Acid Plus Standard-Exposure CsA in Renal-Transplant Recipients. Am J Transplant 2010; 10: 1401–1413. 2. Cibrik D et al. Randomized Trial of Everolimus-Facilitated Calcineurin Inhibitor Minimization Over 24 Months in Renal Transplantation. Transplantation 2013; 95: 933–942. 3. De Simone P et al. Everolimus With Reduced Tacrolimus Improves Renal Function in De Novo Liver Transplant Recipients: A Randomized Controlled Trial. Am J Transplant 2012; 12: 3008–3020. 4. Saliba F et al. Renal Function at Two Years in Liver Transplant Patients Receiving Everolimus: Results of a Randomized, Multicenter Study. Am J Transplant 2013; 13: 1734–1745. 5. Brennan DC et al. Cytomegalovirus Incidence Between Everolimus Versus Mycophenolate in De Novo Renal Transplants: Pooled Analysis of Three Clinical Trials. Am J Transplant 2011; 11: 2453–2462. 6. Eisen HJ et al. Everolimus Versus Mycophenolate Mofetil in Heart Transplantation: A Randomized, Multicenter Trial. Am J Transplant 2013; 13: 1203–1216. 7. Kobashigawa J et al. Everolimus is associated with a reduced incidence of cytomegalovirus infection following de novo cardiac transplantation. Transpl Infect Dis 2013; 15: 150–162. 8. Fachinformation CERTICAN.

Das Novartis-Portfolio.

